Asthma care plan for

CONFIDENTIAL: Staff are trained in asthma first aid (see overleaf) and can provide routine asthma medication as authorised in this care plan by the treating doctor. Please advise staff in writing of any changes to this plan. To be completed by the treating doctor and parent/guardian, for supervising staff and emergency medical personnel. PLEASE PRINT CLEARLY						Photo of child Plan date / /202	
						Review date	
Child's name		Date of birth				/202	
Managing an asthma attack Staff are trained in asthma first aid (see over	leaf). Please write down anyt	hing different th	iis child mi	ght need if th	ney have an	asthma attack:	
Daily asthma management				1			
This child's usual asthma signs Cough Wheeze Difficulty breathing Other (please describe)						child's asthma (eg ke) — please detail:	
Does this child usually tell an adult if s/he is Does this child need help to take asthma me Does this child use a mask with a spacer? *Does this child need a blue reliever puffer	edication?	☐ Yes ☐ Yes ☐ Yes ☐ Yes ☐ Yes	_ N	lo lo lo			
Medication plan If this child needs asthma medication, please	e detail below and make sure	e the medication	and space	er/mask are su	upplied to st	aff.	
Name of medication and colour	Dose/numl	per of puffs				Time required	
	-				,		
Doctor Name of doctor Address	Parent/Guardian I have read, understood and agreed with this care plan and any attachments listed. I approve the release of this information to staff and emergency medical personnel. I will notify the staff in writing if there are any changes to these instructions. I understand staff will seek emergency medical help as needed and that I am responsible for payment of any emergency medical costs. Emergency Contact name				nformation		
Phone	Signature	Signature Date		Mobile			



Name

Signature





Email

Asthma First Aid

- **1** Sit the person upright
 - Be calm and reassuring
 - Do not leave them alone



- Give 4 separate puffs of blue/grey reliever puffer
 - Shake puffer
 - Put 1 puff into spacer
 - Take 4 breaths from spacer

Repeat until 4 puffs have been taken

Remember: Shake, 1 puff, 4 breaths

OR give 2 separate doses of a Bricanyl inhaler (age 6 & over) or a Symbicort inhaler (over 12)



- Wait 4 minutes
 - If there is no improvement, give <u>4 more separate puffs of blue/grey reliever</u> as above

OR give 1 more dose of Bricanyl or Symbicort inhaler



- If there is still no improvement call emergency assistance Dial Triple Zero (000)
 - Say 'ambulance' and that someone is having an asthma attack
 - Keep giving <u>4 separate puffs</u> every <u>4 minutes</u> until emergency assistance arrives

OR give 1 dose of a Bricanyl or Symbicort every 4 minutes - up to 3 more doses of Symbicort



- Call emergency assistance immediately Dial Triple Zero (000)
- If the person is not breathing
- If the person's asthma suddenly becomes worse or is not improving
- If the person is having an asthma attack and a reliever is not available
- If you are not sure if it's asthma
- If the person is known to have Anaphylaxis follow their Anaphylaxis Action Plan, then give Asthma First Aid

Blue/grey reliever medication is unlikely to harm, even if the person does not have asthma



Contact your local Asthma Foundation

1800 ASTHMA Helpline (1800 278 462) asthmaaustralia.org.au